



#4
COPY OF PAPERS
ORIGINALLY FILED



Kongeriget Danmark

Patent application No.: PA 2001 00270

Date of filing: 17 February 2001

Applicant: Novozymes A/S
Krogshøjvej 36
DK-2880 Bagsværd

This is to certify the correctness of the following information:

The attached photocopy is a true copy of the following document:

- The specification, claims and abstract as filed with the application on the filing date indicated above.



Patent- og
Varemærkestyrelsen
Erhvervsministeriet



Taastrup 01 February 2002


Karin Schlichting
Head of office

17 FEB. 2001

Modtaget

1

REDUCTION OF MALODOR FROM LAUNDRY

FIELD OF THE INVENTION

The invention relates to enzymatic reduction of malodor from laundry.

5

BACKGROUND

Clothes that have been used for e.g. sports activities or in other ways have been exposed to sweat from the wearer during use are often difficult to clean in terms of the smell of sweat and other body odors (malodor), when subjected to wash in a
10 laundry machine.

SUMMARY OF THE INVENTION

We have found that an enzyme having lysostaphin activity is capable of reducing malodor from laundry. Accordingly, the invention provides a method
15 comprising contacting laundry with an enzyme having lysostaphin activity.

In a second aspect, there is provided a composition comprising a surfactant and an enzyme having lysostaphin activity.

In another aspect, an enzyme is used for reducing malodor in laundry.

20 DETAILED DESCRIPTION

Lysostaphin

The enzyme Lysostaphin is a glycyl-glycine endopeptidase from the enzyme class E.C. 3.4.24.75. It hydrolyses the -Gly-Gly- bond in the polyglycine inter-peptide link joining staphylococcal cell wall peptidoglycans. Lysostaphin is commercially
25 available from several suppliers, such as from Sigma-Aldrich, Inc.

An enzyme having lysostaphin activity may be a natural or synthetic variant of lysostaphin, wherein amino acid substitutions or deletions have been introduced. It may also be an amino acid fragment with lysostaphin activity, which is optionally fused to one or more other proteins.

30

Lysostaphin Activity

One unit of lysostaphin activity will reduce the turbidity (absorbance at 620 nm) of a suspension of *Staphylococcus aureus* (ATCC 6538) cells by 50%, when the

initial absorbance is approximately 0.250, after 10 minutes at pH 7.5 and 37 degrees celcius.

Malodor

5 During wear, fabrics are contaminated with microorganisms from the human skin together with sebaceous lipids, sweat and dead skin cells serving as microbial substrates. Particularly at mild wash conditions these microorganisms survive laundering. A consequence of microorganisms surviving household laundering is malodor generation in the fabrics. We have now found that these microorganisms are
10 mainly *Staphylococcus* species.

The malodor of the method of the invention thus comprises all body odors present in laundry originating from contact with the human skin. In an embodiment the malodor may be axillary odors, such as the smell of sweat.

In another embodiment, the malodor may originate from activity of
15 *Staphylococcus* species (such as *S. aureus*, *S. epidermis*, *S. intermedius*, *S. saprophyticus* and *S. hyicus*).

In yet another embodiment the malodor originates from fabrics which have been in contact with the axilla.

20 Evaluation of Malodor

Wet laundry swatches are placed in tinted 200 ml glasses with snap lids. A trained sensory panel (10 persons) evaluates the odor by sniffing the headspace over the wet items and indicating the total odor intensity. The odor intensity is indicated on a scale from 0 to 15 where 0 equals 'no malodor' and 15 equals 'very
25 strong malodor'. All evaluations are performed twice (double determinations). The swatches are evaluated after approximately 24 hours and after 48 hours (swatches are kept in the glasses at all times). An average of all 20 evaluations (10 persons each evaluate twice) is calculated after both 24 hours and 48 hours. These average values are referred to as "Malodor index (24 hours)" and "Malodor index (48 hours)".

30

Laundry

The laundry of the method of the invention comprises all kinds of textile items or fabrics suitable for being used as clothes or for personal use in other ways comprising contact with the human skin.

5 Methods and Uses

By contacting laundry with an enzyme having lysostaphin activity, as defined in the method of the invention, the "Malodor index (48 hours)", as defined above, may be reduced by at least 10% (preferably 20%, more preferably 30%, most preferably 40%, and in particular 50%) compared to laundry which has not been contacted with
10 an enzyme having lysostaphin activity.

The method of the invention may also result in killing or inhibiting growth of microbial cells in laundry. In an embodiment the microbial cells are bacteria, such as *Staphylococcus* species. In another embodiment, the method of the invention may result in a reduction in the number of living microbial cells of at least 25%, preferably
15 at least 50%, more preferably at least 90%, and most preferably at least 99%.

In the context of the present invention the term "inhibiting growth of microbial cells" is intended to mean that the cells are in the non-growing state, i.e., that they are not propagating. The term "microbial cells" denotes bacterial cells (such as *Staphylococcus* species), fungal cells or algae, and the term "microorganism"
20 denotes a fungus (including yeasts) or a bacterium

The present invention covers use of an enzyme for reducing malodor from laundry items. In an embodiment, the enzyme may have lysostaphin activity. The invention may also be used for reducing the number of living bacteria in laundry; for reducing allergens in laundry; or for sterilizing laundry.
25

Detergent composition

Lysostaphin may be added to and thus become a component of a detergent composition.

The detergent composition of the invention may for example be formulated as a
30 hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general

household hard surface cleaning operations, or be formulated for hand or machine dishwashing operations.

In a specific aspect, the invention provides a detergent additive comprising lysostaphin. The detergent additive as well as the detergent composition may
5 comprise one or more other enzymes such as a protease, a lipase, a cutinase, an amylase, a carbohydrase, a cellulase, a pectinase, a mannanase, an arabinase, a galactanase, a xylanase, an oxidase, e.g., a laccase, and/or a peroxidase.

In general the properties of the chosen enzyme(s) should be compatible with the selected detergent, (i.e. pH-optimum, compatibility with other enzymatic and non-
10 enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

Proteases: Suitable proteases include those of animal, vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. The protease may be a serine protease or a metallo protease,
15 preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from *Bacillus*, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270 and
20 WO 94/25583.

Examples of useful proteases are the variants described in WO 92/19729, WO 98/20115, WO 98/20116, and WO 98/34946, especially the variants with substitutions in one or more of the following positions: 27, 36, 57, 76, 87, 97, 101, 104, 120, 123, 167, 170, 194, 206, 218, 222, 224, 235 and 274.

25 Preferred commercially available protease enzymes include Alcalase™, Savinase™, Primase™, Everlase™, Esperase™, and Kannase™ (Novozymes A/S), Maxatase™, Maxacal™, Maxapem™, Properase™, Purafect™, Purafect OxP™, FN2™, and FN3™ (Genencor International Inc.).

Lipases: Suitable lipases include those of bacterial or fungal origin. Chemically
30 modified or protein engineered mutants are included. Examples of useful lipases include lipases from *Humicola* (synonym *Thermomyces*), e.g. from *H. lanuginosa* (*T. lanuginosus*) as described in EP 258 068 and EP 305 216 or from *H. insolens* as described in WO 96/13580, a *Pseudomonas* lipase, e.g. from *P. alcaligenes* or *P.*

pseudoalcaligenes (EP 218 272), *P. cepacia* (EP 331 376), *P. stutzeri* (GB 1,372,034), *P. fluorescens*, *Pseudomonas* sp. strain SD 705 (WO 95/06720 and WO 96/27002), *P. wisconsinensis* (WO 96/12012), a *Bacillus* lipase, e.g. from *B. subtilis* (Dartois et al. (1993), *Biochemica et Biophysica Acta*, 1131, 253-360), *B.*
5 *stearothermophilus* (JP 64/744992) or *B. pumilus* (WO 91/16422).

Other examples are lipase variants such as those described in WO 92/05249, WO 94/01541, EP 407 225, EP 260 105, WO 95/35381, WO 96/00292, WO 95/30744, WO 94/25578, WO 95/14783, WO 95/22615, WO 97/04079 and WO 97/07202.

10 Preferred commercially available lipase enzymes include Lipolase™, Lipolase Ultra™ and Lipoprime™ (Novozymes A/S).

Amylases: Suitable amylases (α and/or β) include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, α -amylases obtained from *Bacillus*, e.g. a special strain of *B.*
15 *licheniformis*, described in more detail in GB 1,296,839.

Examples of useful amylases are the variants described in WO 94/02597, WO 94/18314, WO 96/23873, and WO 97/43424, especially the variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 181, 188, 190, 197, 202, 208, 209, 243, 264, 304, 305, 391, 408, and
20 444.

Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™ and BAN™ (Novozymes A/S), Rapidase™ and Purastar™ (Genencor International Inc.).

Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases
25 include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g. the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in US 4,435,307, US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259.

Especially suitable cellulases are the alkaline or neutral cellulases having colour
30 care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372, WO 96/11262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, US

5,457,046, US 5,686,593, US 5,763,254, WO 95/24471, WO 98/12307 and PCT/DK98/00299.

Commercially available cellulases include Celluzyme™, and Carezyme™ (Novozymes A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.),
5 and KAC-500(B)™ (Kao Corporation).

Peroxidases/Oxidases: Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g. from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO
10 95/10602, and WO 98/15257.

The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these enzymes. A detergent additive of the invention, i.e. a separate additive or a combined additive, can be formulated e.g. as a granulate, a
15 liquid, a slurry, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol,
20 PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed
25 techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

The detergent composition of the invention may be in any convenient form, e.g.,
30 a bar, a tablet, a powder, a granule, a paste or a liquid. A liquid detergent may be aqueous, typically containing up to 70 % water and 0-30 % organic solvent, or non-aqueous.

The detergent composition comprises one or more surfactants, which may be non-ionic including semi-polar and/or anionic and/or cationic and/or zwitterionic. The surfactants are typically present at a level of from 0.1% to 60% by weight.

When included therein the detergent will usually contain from about 1% to about 5 40% of an anionic surfactant such as linear alkylbenzenesulfonate, alpha-olefinsulfonate, alkyl sulfate (fatty alcohol sulfate), alcohol ethoxysulfate, secondary alkanesulfonate, alpha-sulfo fatty acid methyl ester, alkyl- or alkenylsuccinic acid or soap.

When included therein the detergent will usually contain from about 0.2% to 10 about 40% of a non-ionic surfactant such as alcohol ethoxylate, nonylphenol ethoxylate, alkylpolyglycoside, alkyldimethylamineoxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, polyhydroxy alkyl fatty acid amide, or N-acyl N-alkyl derivatives of glucosamine ("glucamides").

The detergent may contain 0-65 % of a detergent builder or complexing agent 15 such as zeolite, diphosphate, triphosphate, phosphonate, carbonate, citrate, nitrilotriacetic acid, ethylenediaminetetraacetic acid, diethylenetriaminepentaacetic acid, alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates (e.g. SKS-6 from Hoechst).

The detergent may comprise one or more polymers. Examples are 20 carboxymethylcellulose, poly(vinylpyrrolidone), poly(ethylene glycol), poly(vinyl alcohol), poly(vinylpyridine-N-oxide), poly(vinylimidazole), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

The detergent may contain a bleaching system which may comprise a H_2O_2 25 source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetylenediamine or nonanoyloxybenzenesulfonate. Alternatively, the bleaching system may comprise peroxyacids of e.g. the amide, imide, or sulfone type.

The enzyme(s) of the detergent composition of the invention may be stabilized 30 using conventional stabilizing agents, e.g., a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e.g., an aromatic borate ester, or a phenyl boronic acid derivative such as 4-

formylphenyl boronic acid, and the composition may be formulated as described in e.g. WO 92/19709 and WO 92/19708.

The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bactericides, optical brighteners, hydrotropes, tarnish inhibitors, or perfumes.

It is at present contemplated that in the detergent compositions any enzyme, in particular lysostaphin, may be added in an amount corresponding to 0.01-100 mg of enzyme protein per liter of wash liquor, preferably 0.05-10 mg of enzyme protein per liter of wash liquor, more preferably 0.1-5 mg of enzyme protein per liter of wash liquor, and most preferably 0.1-1 mg of enzyme protein per liter of wash liquor.

Lysostaphin may additionally be incorporated in the detergent formulations disclosed in WO 97/07202 which is hereby incorporated as reference.

The present invention is further illustrated in the following examples, which are not in any way intended to limit the scope of the invention as claimed.

EXAMPLES

The chemicals used in the following examples were commercial products of at least reagent grade.

The Malthus Flexi M2060 instrument is available from Malthus Instruments Limited, England.

Tryptone Soya Broth (TSB) is available from Oxoid, England.

CFU: Colony Forming Units.

EXAMPLE 1

Evaluation of odour in swatches soiled with human axillary sweat and sebum

The odour reducing effect of lysostaphin is tested in a one-cycle washing trial carried out in a thermostated Terg-O-tometer (available from United States Testing Co, 1415 Park Ave, Hoboken NJ07030).

Experimental conditions:

Soiling

Swatches of 100% polyester or 100% cotton (10 × 14 cm, previously cleaned by solvent extraction - hexane for polyester, chloroform for cotton - using a Soxhlet) are
5 soiled by applying human male axillary sweat and sebum from the armpits and upper body of male runners after extensive exercise. Two swatches (10 × 14 cm) are used for each male - one swatch is made of 100% polyester and one is made of 100% cotton. The left armpit and left part of the upper body is wiped with one swatch, and the right armpit and right part of the upper body is wiped with the other swatch after
10 exercising. This procedure is performed twice before washing. Prior to the washing, each swatch is cut into 12 equally sized pieces and distributed between four Terg-O-tometer wash beakers. Cotton and polyester textiles are kept apart.

Washing

- 15 Washing is done in Terg-O-tometer using phosphate buffer (0.05 M, pH 7.5).
Temperature: 32°C
Lysostaphin dose: 5 mg/L (Sigma L4402)
Wash time: 12 minutes
Wash liquid: 1000 mL per wash beaker
20 Rinsing: 15 minutes in running tap water

Sensory evaluation

After wash the wet swatches are evaluated and "Malodor index (24 hours)" and "Malodor index (48 hours)" are calculated. The results show that a lower "Malodor
25 index (48 hours)" is obtained with the swatches that has been washed with lysostaphin compared to those that has not been washed with lysostaphin.

EXAMPLE 2

Removal of bacteria from textile by lysostaphin

- 30 *Staphylococcus aureus* (ATCC 6538) was grown overnight (TSB; Tryptone Soya Broth) and inoculated to approximately 10^3 Colony forming units/ml (CFU/ml) in diluted TSB (1:1). Sterile cotton swatches were inoculated overnight in the diluted TSB allowing *S.aureus* to grow on the textile. The swatches were rinsed in sterile

water for 30 seconds and dried in sterile air for 30 minutes. Swatches were washed in a beaker with stirring at 32°C for 20 minutes in either phosphate buffer (0.05 M, pH 7.5) or in a liquid U.S. detergent of a commercial type (0.75 g/liter) with and without addition of 5 mg/L Lysostaphin (Sigma L4402). Three swatches were washed in each beaker. After wash all swatches were rinsed in sterile water for 10 minutes and dried in sterile air. The number of living *S. aureus* on the swatches was determined by incubation of the swatches in Malthus tubes with CASO medium (Merck 1.05459). The bactericidal activity was determined by incubation in Malthus. The detection times (dt) measured by the Malthus instrument were converted to CFU/swatch by a calibration curve (Johansen et al. 1999, Methods in Enzymology, vol 310, p. 353-360). Direct Malthus measurements were used when enumerating total survival cells. By the direct measurements, the cell metabolism was determined by conductance measurements in the growth substrate. The swatches were after enzyme treatment transferred to the Malthus cell. As cells attached to the textile are growing, the cell metabolism will change the conductance in the growth medium. When the conductance change is measurable by the Malthus, a detection time (dt) will be recorded. The dt's were converted to colony counts by use of a calibration curve relating CFU/swatch to dt.

20 Results:

Swatch	Beaker	Average number of <i>S. aureus</i> on each swatch (three swatches washed in each beaker)
Reference swatches before wash		8.8×10^7 CFU/swatch
Washed in phosphate buffer	1	2.2×10^7 CFU/swatch
	2	7.3×10^6 CFU/swatch
Washed in phosphate buffer with lysostaphin (5 mg/L)	1	4.6×10^6 CFU/swatch
	2	1.2×10^5 CFU/swatch
	3	4.8×10^5 CFU/swatch

Lysostaphin resulted in a reduction of the microbial cell number of approximately 10^1 to 10^2 CFU/swatch. A reduction in cell number was also determined after lysostaphin

treatment in detergent, however, the determination of the exact cell number was not possible by using the Malthus. But a delay in outgrowth of the microorganism was observed visually from the swatches washed in detergent with lysostaphin, this delay corresponds to a lower cell number on the swatches compared to the swatches washed without lysostaphin.

CLAIMS

1. A method comprising contacting laundry with an enzyme having lysostaphin activity.

5

2. The method of claim 1, wherein the enzyme is lysostaphin.

3. The method of claim 1, which is a method for reducing malodor from laundry.

10 4. The method of claim 3, wherein the "malodor index (48 hours)" is reduced 10% compared to laundry which has not been contacted with an enzyme having lysostaphin activity.

5. A composition comprising a surfactant and an enzyme having lysostaphin activity.

15

6. The composition of claim 3, wherein the enzyme is lysostaphin.

7. Use of an enzyme for reducing malodor from laundry.

20 8. Use of an enzyme having lysostaphin activity for reducing malodor from laundry.

9. Use of lysostaphin for reducing malodor from laundry.

ABSTRACT

Title: Reduction of Malodor from Laundry

- 5 We have found that an enzyme having lysostaphin activity is capable of reducing malodor from laundry. Accordingly, there is provided a method comprising contacting laundry with an enzyme having lysostaphin activity; and a composition comprising a surfactant and an enzyme having lysostaphin activity.